

ADDITION THEOREMS
from the
STANDPOINT OF ABEL'S THEOREM.

submitted by
WALTER REYNOLDS, JR.

as thesis for
DEGREE OF MASTER OF SCIENCE
from
GEORGIA SCHOOL OF TECHNOLOGY, JUNE 4, 1934.

35003

APPROVED BY: _____ .

APPROVED BY: _____ .

APPROVED BY: *[Signature]* .

INTRODUCTION

In this paper there is shown in detail the proofs of certain addition theorems from the standpoint of Abel's Theorem.

ABELIAN INTEGRALS

Any integral $I = \int R(xy)dx$ where $R(xy)$ is a rational function of x and y , where y is an algebraic function defined by an equation $F(xy) = 0$ is called an Abelian Integral attached to that curve. To obtain the determination of this integral it is necessary to assign a lower limit x_0 and a corresponding value y_0 is chosen among roots of equation $F(x_0, y) = 0$. On going from point x_0 to any point x by all possible paths, all values of the integral I are included in one of the formulas $I = I_K + m_1 w_1 + m_2 w_2 + \dots - m_n w_n$ ($K = 1, 2, 3, \dots, n$) where I_1, I_2, \dots, I_n are values of the integral which correspond to certain definite paths; m_1, m_2, \dots, m_n , and w_1, w_2, \dots, w_n , are periods. These periods are of two kinds; one kind results from loops described about the poles of $F(xy) = 0$; these are Polar Periods. The other comes from closed paths surrounding several critical points, called Cycles; these are called Cyclic Periods. The number of distinct Cyclic periods depends only on the algebraic function considered $F(xy) = 0$; it is equal to $2p$, where p denotes the deficiency of the curve. From point of view of singularities 3 classes of Abelian Integrals are distinguished.

(1) Those which remain finite in the neighborhood of every value of x are called First Kind.

(2) Those which have a single pole are called the Second Kind.

(3) Those having two logarithmic singular points are called Third Kind.

ABEL'S THEOREM

Consider a plane curve $F(xy) = 0$ and let $\phi(xy) = 0$ be the equation of another plane algebraic curve. These curves have n points in common $(x_1, y_1), (x_2, y_2), \dots$ the number n equaling the product of degrees of the two curves. Let $R(xy)$ be a rational function and consider the following sum

(1) $I = \sum_{i=1}^n \int_{x_0 y_0}^{x_i y_i} R(xy) dx$ where $\int_{x_0 y_0}^{x_i y_i} R(xy) dx$ denotes the Abelian integral taken from a fixed point x_0 to a point x_i along a path which leads y from an initial value y_0 to final of y_i , the initial value y_0 being same for all these integrals.

Suppose, now, some of the coefficients a, b, c, \dots of $\phi(xy) = 0$ are variable. When these coefficients vary continuously the points x_i vary continuously, and if none of these pass thru a discontinuity of $\int R(xy) dx$, the sum I itself varies continuously, provided we follow the continuous variation of each of the integrals contained in it along the entire path described by corresponding upper limit. The sum I is therefore a function of the parameters a, b, c, \dots , whose analytic form is considered next.

Denoting the total differential of any function U by dU , with respect to variables a, b, c, \dots , we have,

$dU = \frac{\partial U}{\partial a} da + \frac{\partial U}{\partial b} db + \dots$. By (1) $dI = \sum_{i=1}^n R(x_i, y_i) dx_i$ and from the two relations $F(x_i, y_i) = 0$ and $\phi(x_i, y_i) = 0$ there is derived

$$\frac{\partial F}{\partial x_i} dx_i + \frac{\partial F}{\partial y_i} dy_i = 0$$

$$\text{and} \quad \frac{\partial \phi}{\partial x_i} dx_i + \frac{\partial \phi}{\partial y_i} dy_i + d\phi_i = 0$$

and, consequently $dx_i = \psi(x_i, y_i) d\phi_i$ where $\psi(x_i, y_i)$ is a rational function of x_i, y_i, a, b, c, \dots and where ϕ_i stands for $\phi(x_i, y_i)$, we obtain,

$$dI = \sum_{i=1}^n R(x_i, y_i) \psi(x_i, y_i) d\phi_i$$

The coefficient of da is a rational symmetric function of the coordinates of n points (x_i, y_i) common to the two curves $F(xy) = 0$ and $\phi(xy) = 0$. The theory of elimination tells us that this function is a rational function of coefficients of the two polynomials $F(xy) = 0$ and $\phi(xy) = 0$ and consequently a rational function of a, b, c, \dots . Evidently the same is true of the coefficients of db, dc, \dots and I will be obtained by integration of a total differential

$I = \int (f_1 da + f_2 db + f_3 dc) \dots$ where f_1, f_2, f_3, \dots are rational functions of a, b, c, \dots and the integration cannot introduce any other transcendental other than logarithms.

The sum I is therefore equal to a Rational Function of the Coefficients a, b, c, \dots plus a sum of logarithms

of rational functions of the coefficients a, b, c, \dots , each of these logarithms being multiplied by a constant factor.

The above is a statement of Abel's Theorem in its most general form.

In geometric language this can be stated as follows: The sum of the values of an Abelian Integral, taken from a common origin to the n points of intersection of the given curve with a variable curve of degree m , $\phi(xy) = 0$, is equal to a rational function of the coefficients of $\phi(xy) = 0$ plus a sum of a finite number of logarithms of rational functions of the same coefficients, each logarithm being multiplied by a constant factor.

Abel's Theorem has a definite meaning only if paths described by $x_1, x_2, x_3, \dots, x_n$ are taken into account.

The theorem becomes simpler when the Abelian integral is of the first kind. If $f_1, f_2, f_3, \dots, f_k$ were not identically zero, it would be possible to find a system of values $a = a', b = b', c = c', \dots$ for which I would become infinite. Let $(x'_1, y'_1) \dots (x'_n, y'_n)$ be the intersections of the curves $F(xy)$ and $\phi(xy)$ which correspond to the parametric values a', b', c' .

Then $\int_{x_0, y_0}^{xy} R(xy) dx$ becomes infinite when the upper limit approaches one of the points (x'_i, y'_i) ; Since this is impossible when the Integral is of the first kind, we have $dI = 0$, and, consequently $I = a$ constant, as a, b, c , vary

continuously.

Abel's Theorem can then be stated as follows: Given a fixed curve $F(xy) = 0$ and a variable curve $\phi(xy) = 0$ of degree m , the sum of the increments of an Abelian Integral of the first kind attached to the curve $F(xy)$ along the continuous curves described by the points of intersection of $F(xy) = 0$ with $\phi(xy) = 0$ is equal to zero.

Note: References for theory are made especially to the following three texts which were consulted:

Goursat's: Mathematical Analysis Vol.2 Part1


H.F. Baker's: Abelian Functions.

Harkness and Morley: Theory of Functions.

SECTION (1)

THE ADDITION THEOREM FOR $\log x = \int_1^x \frac{dx}{x}$

- (1) Let $y = \frac{1}{x}$ represent the fixed curve
 (2) And $y = x^2 + ax + b$ the variable curve



$$U(ab) = \int_1^{x_1(ab)} \frac{dx}{x} + \int_1^{x_2(ab)} \frac{dx}{x} + \int_1^{x_3(ab)} \frac{dx}{x} \quad \text{by Abel's theorem}$$

$$(3) \quad U_a = y_1 x_{1a} + y_2 x_{2a} + y_3 x_{3a}$$

On eliminating y from (1) and (2) we have,

$$(4) \quad \phi(abx) = x^3 + ax^2 + bx - 1 \quad \text{whose roots are the abscissas of the intersections of (1) and (2)}$$

$$\text{Now } \phi'_x \cdot x_a + \phi_a = \phi'_x \cdot x_a + x^2 = 0$$

$$\text{Therefore } x_a = -\frac{x^2}{\phi'_x}$$

Substitution in equation (3) gives

$$(5) \quad U_a = \frac{1}{x_1} \cdot \frac{-x_1^2}{\phi'_{x_1}} + \frac{1}{x_2} \cdot \frac{-x_2^2}{\phi'_{x_2}} + \frac{1}{x_3} \cdot \frac{-x_3^2}{\phi'_{x_3}}$$

$$\text{Or } U_a = -\left[\frac{x_1}{\phi'_{x_1}} + \frac{x_2}{\phi'_{x_2}} + \frac{x_3}{\phi'_{x_3}} \right]$$

where, $\phi'_{x_1} = 3x_1^2 + 2ax_1 + b$
 $\phi'_{x_2} = 3x_2^2 + 2ax_2 + b$
 $\phi'_{x_3} = 3x_3^2 + 2ax_3 + b$

$$\text{Therefore } U_a = - \left[\frac{x_1}{3x_1^2 + 2ax_1 + b} + \frac{x_2}{3x_2^2 + 2ax_2 + b} + \frac{x_3}{3x_3^2 + 2ax_3 + b} \right]$$

(6) By elementary operations we have as the numerator of U_a ,

$$- [9x_1 x_2 x_3 (x_2 x_3 + x_1 x_3 + x_1 x_2) + 6ax_1 x_2 x_3 (x_1 + x_2 + x_3) + 3bx_1 (x_3^2 + x_1 x_2 + x_1 x_3) + 6ax_1 x_2 x_3 (x_2 + x_3 + x_1) + 4a^2 x_1 x_2 x_3 (1+1+1) + 2abx_1 (x_2 + x_2 + x_3) + 3bx_2 (x_1 x_2 + x_3^2 + x_2 x_3) + 2abx_2 (x_1 + x_3 + x_3) + b^2 (x_1 + x_2 + x_3)]$$

We now simplify this expression as follows:

From (4) we have,

$$(7) \quad \begin{cases} x_1 + x_2 + x_3 = -a \\ x_1 x_2 + x_1 x_3 + x_2 x_3 = b \\ x_1 x_2 x_3 = 1 \end{cases}$$

Substituting relations from (7) in (6) we get,

$$- [9b - 6a^2 + 3bx_1 (x_3^2 + x_1 x_2 + x_1 x_3) - 6a^2 + 12a^2 + 4ab^2 + 3bx_2 (x_1 x_2 + x_3^2 + x_2 x_3) - ab^2] =$$

$$(8) \quad - [9b + 3ab^2 + 3b(x_1 x_3^2 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + x_2 x_3^2 + x_2^2 x_3)]$$

From (7) $(x_1 x_2 + x_1 x_3 + x_2 x_3)(x_1 + x_2 + x_3) = x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + x_2^2 x_3 + x_1 x_3^2 + x_2 x_3^2 + 3x_1 x_2 x_3$

Therefore $(b)(-a) = x_1 x_3^2 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + x_2 x_3^2 + x_2^2 x_3 + 3$ or $x_1 x_3^2 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + x_2 x_3^2 + x_2^2 x_3 = -ab - 3$

Substitution in (8) gives for the numerator of U_a ,
 $- [9b + 3ab^2 + 3b(-ab - 3)] = - [9b + 3ab^2 - 3ab^2 - 9b] = 0$

$$\text{Hence } U_a = 0$$

Likewise, for U_b , we have,

$$(9) \quad U_b = y_1 x_{1b} + y_2 x_{2b} + y_3 x_{3b}$$

and $\phi'_x \cdot x_b + \phi_b = \phi'_x \cdot x_b + x = 0$

Therefore $x_b = - \frac{x}{\phi'_x}$

Substitution in (9) gives,

$$U_b = \frac{1}{x_1} \cdot \frac{-x_1}{\phi'_{x_1}} + \frac{1}{x_2} \cdot \frac{-x_2}{\phi'_{x_2}} + \frac{1}{x_3} \cdot \frac{-x_3}{\phi'_{x_3}}$$

Or

$$U_b = - \left[\frac{1}{\phi'_{x_1}} + \frac{1}{\phi'_{x_2}} + \frac{1}{\phi'_{x_3}} \right]$$

$$(10) \quad U_b = - \left[\frac{1}{3x_1^2 + 2ax_1 + b} + \frac{1}{3x_2^2 + 2ax_2 + b} + \frac{1}{3x_3^2 + 2ax_3 + b} \right]$$

By elementary reductions the numerator of (10) becomes,

$$(11) \quad - [9(x_1^2 x_2^2 + x_1^2 x_3^2 + x_2^2 x_3^2) + 6a(x_2 x_3^2 + x_1 x_3^2 + x_3 x_1^2 + x_2 x_1^2 + x_3 x_2^2 + x_1 x_2^2) + 6b(x_1^2 + x_2^2 + x_3^2) + 4a^2(x_2 x_3 + x_1 x_3 + x_1 x_2) + 4ab(x_1 + x_2 + x_3) + 3b^2]$$

Squaring the first equation of (7) and transposing, we obtain:

$$x_1^2 + x_2^2 + x_3^2 = a^2 - 2b$$

And from U_ω it has been shown that,

$$x_1 x_3^2 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + x_2 x_3^2 + x_2^2 x_3 = -ab - 3$$

Also, from the second equation in (7)

$$x_1^2 x_2^2 + x_2^2 x_3^2 + x_1^2 x_3^2 = b^2 - 2x_1 x_2 x_3 (x_1 + x_2 + x_3)$$

Therefore

$$x_1^2 x_2^2 + x_2^2 x_3^2 + x_1^2 x_3^2 = b^2 + 2a$$

Hence, substituting relations from cubic in (11) we obtain; for the numerator of U_b ,

$$- [9(b^2 + 2a) + 6a(-ab - 3) + 6b(a^2 - 2b) + 4a^2(b) + 4ab(-a) + 3b^2] = - [9b^2 + 18a - 6a^2b - 18a + 6a^2b - 12b^2 + 4a^2b - 4a^2b + 3b^2] = 0$$

Therefore $U_b = 0$

Hence, $U_\omega = 0$, $U_b = 0$, and consequently

$$U(ab) = \text{a constant} = C$$

And

$$\int_1^{x_1} \frac{dx}{x} + \int_1^{x_2} \frac{dx}{x} + \int_1^{x_3} \frac{dx}{x} = C$$

As no restrictions were placed on a and b we take

$a = -3$, $b = 3$ The intersection then becomes,

$$x^3 - 3x^2 + 3x - 1 = 0 \text{ with a triple root } x_1 = x_2 = x_3 = 1$$

Therefore we have, $\int_1^{x_1} \frac{dx}{x} + \int_1^{x_2} \frac{dx}{x} + \int_1^{x_3} \frac{dx}{x} = C$

Hence $0 = C$

$$\text{And } \int_1^{x_1} \frac{dx}{x} + \int_1^{x_2} \frac{dx}{x} + \int_1^{x_3} \frac{dx}{x} = 0$$

$$(12) \text{ Transposing, } \int_1^{x_1} \frac{dx}{x} + \int_1^{x_2} \frac{dx}{x} = -\int_1^{x_3} \frac{dx}{x} = \int_1^{\frac{1}{x_3}} \frac{dx}{x}$$

As $x_1 x_2 x_3 = 1$ we get $x_3 = \frac{1}{x_1 x_2}$, therefore $\frac{1}{x_3} = x_1 x_2$


Hence, substituting in (12) we get,

$$\int_1^{x_1} \frac{dx}{x} + \int_1^{x_2} \frac{dx}{x} = \int_1^{x_1 x_2} \frac{dx}{x}$$

Or, $\log x_1 + \log x_2 = \log x_1 x_2$, which is

THE ADDITION THEOREM FOR LOGARITHMS


The curves below show how the intersection occurs for the special case chosen, $a = -3$, $b = 3$



SECTION (2)

THE ADDITION THEOREM FOR ARC TANGENT $x = \int_0^x \frac{dx}{x^2 + 1}$

- (1) Let $y = \frac{1}{x^2 + 1}$ represent the fixed curve
 (2) And $y = ax + b$ the variable curve



$$U(ab) = \int_0^{x_1(ab)} \frac{dx}{x^2 + 1} + \int_0^{x_2(ab)} \frac{dx}{x^2 + 1} + \int_0^{x_3(ab)} \frac{dx}{x^2 + 1}$$

On eliminating y between (1) and (2) we get,

- (3) $ax^3 + bx^2 + ax + b - 1 = 0$ whose roots are the abscissas of the intersections of (1) and (2)

Proceeding as in Section (1), we have

(4) $U_a = y_1 x_{1a} + y_2 x_{2a} + y_3 x_{3a}$

Since $\phi(abx) = ax^3 + bx^2 + ax + b - 1$

Therefore $\phi'_x = 3ax^2 + 2bx + a$

Here $\phi'_x \cdot x_a + \phi_a = \phi'_x \cdot x_a + (x^3 + x) = 0$

Hence $x_a = -\frac{x^3 + x}{\phi'_x} = -\frac{x(x^2 + 1)}{\phi'_x} = -\frac{x}{y \phi'_x}$

Substitution in (4) gives,

$$U_a = y_1 \cdot \frac{-x_1}{y_1 \phi'_{x_1}} + y_2 \cdot \frac{-x_2}{y_2 \phi'_{x_2}} + y_3 \cdot \frac{-x_3}{y_3 \phi'_{x_3}}$$

$$U_a = - \left[\frac{x_1}{3ax_1^2 + 2bx_1 + a} + \frac{x_2}{3ax_2^2 + 2bx_2 + a} + \frac{x_3}{3ax_3^2 + 2bx_3 + a} \right]$$

Hence, we get, as the numerator of U_a ,

$$(5) \quad -[x_1(3ax_2^2 + 2bx_2 + a)(3ax_3^2 + 2bx_3 + a) + x_2(3ax_1^2 + 2bx_1 + a)(3ax_3^2 + 2bx_3 + a) + x_3(3ax_1^2 + 2bx_1 + a)(3ax_2^2 + 2bx_2 + a)]$$

By elementary operations (5) can be written:

$$(6) \quad -[9a^2(x_1x_2x_3)(x_1x_2 + x_1x_3 + x_2x_3) + 6ab(x_1x_2x_3)(x_2 + x_3 + x_2 + x_1 + x_1 + x_3) + 4b^2(x_1x_2x_3 + x_1x_2x_3 + x_1x_2x_3) + 2ab(x_2x_3 + x_2x_3 + x_1x_2 + x_1x_3 + x_1x_2 + x_1x_3) + a^2(x_3 + x_2 + x_1) + 3a^2(x_2^2x_3 + x_2x_3^2 + x_1x_2^2 + x_1^2x_3 + x_1^2x_2 + x_1x_3^2)]$$

From (3) we have,

$$ax^3 + bx^2 + ax + b - 1 = 0, \text{ or } x^3 + \frac{b}{a}x^2 + x + \frac{b-1}{a} = 0$$

Therefore, we obtain,

$$(7) \quad \begin{cases} x_1 + x_2 + x_3 = -\frac{b}{a} \\ x_1x_2 + x_2x_3 + x_1x_3 = 1 \\ x_1x_2x_3 = \frac{1-b}{a} \end{cases}$$

From (7) by multiplying first equation by the second we get,
 $x_2^2x_3 + x_2x_3^2 + x_1x_2^2 + x_1^2x_3 + x_1^2x_2 + x_1x_3^2 = (x_1 + x_2 + x_3)(x_1x_2 + x_1x_3 + x_2x_3) - 3x_1x_2x_3 = -\frac{b}{a} - 3\left(\frac{1-b}{a}\right) = \frac{-b-3+3b}{a} = \frac{-3+2b}{a}$

Substituting these relations in (6) we get, as the numerator of U_a ,

$$- [9a(1-b) + 6b(1-b)\left(-\frac{2b}{a}\right) + 12b^2\left(\frac{1-b}{a}\right) + 4ab - ab - 9a + 6ab] = - [9a - 9ab - \frac{12b^2}{a}(1-b) + \frac{12b^2}{a}(1-b) - 9a + 9ab] = 0$$

$$\text{Therefore } U_a = 0$$

To determine U_b we have,

$$(8) \quad U_b = y_1x_{1b} + y_2x_{2b} + y_3x_{3b}$$

$$\phi(ax) = ax^3 + bx^2 + ax + b - 1$$

Therefore $\phi'_x = 3ax^2 + 2bx + a$, whence

$$\phi'_x \cdot x_b + \phi_b = \phi'_x \cdot x_b + (x^2 + 1) = 0$$

$$\text{Therefore } x_b = -\frac{x^2 + 1}{\phi'_x} = -\frac{1}{y \phi'_x}$$

Substitution in (8) gives,

$$\begin{aligned} U_b &= y_1 \cdot \frac{-1}{y_1 \phi'_x} + y_2 \cdot \frac{-1}{y_2 \phi'_{x_2}} + y_3 \cdot \frac{-1}{y_3 \phi'_{x_3}} = \\ (9) \quad & - \left[\frac{1}{3ax_1^2 + 2bx_1 + a} + \frac{1}{3ax_2^2 + 2bx_2 + a} + \frac{1}{3ax_3^2 + 2bx_3 + a} \right] \end{aligned}$$

Clearing (9) the numerator of U_b becomes,

$$\begin{aligned} (10) \quad & - [(3ax_1^2 + 2bx_1 + a)(3ax_2^2 + 2bx_2 + a) + (3ax_1^2 + 2bx_1 + a) \\ & (3ax_3^2 + 2bx_3 + a) + (3ax_2^2 + 2bx_2 + a)(3ax_3^2 + 2bx_3 + a)] \end{aligned}$$

By elementary reductions (10) can be written:

$$\begin{aligned} (11) \quad & - [9a^2(x_1^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2) + 6ab(x_1x_2^2 + x_1x_3^2 + x_2^2x_3 + \\ & x_1^2x_2 + x_1^2x_3 + x_2^2x_3) + 3a^2(x_2^2 + x_3^2 + x_2^2 + x_1^2 + x_1^2 + x_3^2) + \\ & 4b^2(x_1x_2 + x_1x_3 + x_2x_3) + 2ab(x_2 + x_3 + x_2 + x_1 + x_1 + x_3) + 3a^2] \end{aligned}$$

From (7) we get,

$$\begin{aligned} x_1^2 + x_2^2 + x_3^2 &= \frac{b^2}{a^2} - 2(x_1x_2 + x_1x_3 + x_2x_3) = \frac{b^2}{a^2} - 2 = \frac{b^2 - 2a^2}{a^2} \\ x_1^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2 &= 1 - 2(x_1x_2x_3)(x_1 + x_2 + x_3) = \frac{a^2 + 2b - 2b^2}{a^2} \end{aligned}$$

Substituting these relations, and also the ones under (7) in (11) we get, for the numerator of U_b ,

$$\begin{aligned} & - \left[9a^2 \left(\frac{a^2 + 2b - 2b^2}{a^2} \right) + 6ab \left(\frac{-3 + 2b}{a} \right) + 6a^2 \left(\frac{b^2 - 2a^2}{a^2} \right) + 4b^2 + \right. \\ & \left. 4ab \left(-\frac{b}{a} \right) + 3a^2 \right] = - [9a^2 + 18b + 12b^2 - 18b^2 - 18b + 6b^2 - 12a^2 + \\ & 4b^2 - 4b^2 + 3a^2] = 0, \text{ and consequently } U_b = 0 \end{aligned}$$

$$\text{Hence } U_a = 0, U_b = 0$$

And, consequently, $U(ab) = \text{a constant} = C$

Then, we have (12)
$$\int_0^{x_1} \frac{dx}{x^2+1} + \int_0^{x_2} \frac{dx}{x^2+1} + \int_0^{x_3} \frac{dx}{x^2+1} = C$$

If we let $a = 0$ and $b = 1$ in (3) it reduces to $x^2 = 0$, whose roots are: $x_1 = x_2 = 0$, $x_3 = \infty$

and (12) becomes
$$\int_0^0 \frac{dx}{x^2+1} + \int_0^0 \frac{dx}{x^2+1} + \int_0^\infty \frac{dx}{x^2+1} = C$$

Since the first two integrals vanish, we get

$$\int_0^\infty \frac{dx}{x^2+1} = C = \frac{\pi}{2}. \quad \text{Substituting in (12) we obtain:}$$

$$\int_0^{x_1} \frac{dx}{x^2+1} + \int_0^{x_2} \frac{dx}{x^2+1} + \int_0^{x_3} \frac{dx}{x^2+1} = \frac{\pi}{2}$$

Therefore
$$\int_0^{x_1} \frac{dx}{x^2+1} + \int_0^{x_2} \frac{dx}{x^2+1} = \frac{\pi}{2} - \int_0^{x_3} \frac{dx}{x^2+1}$$

Since $\frac{\pi}{2} - \tan^{-1} x_3 = \tan^{-1} \frac{1}{x_3}$, hence we have

$$(13) \quad \int_0^{x_1} \frac{dx}{x^2+1} + \int_0^{x_2} \frac{dx}{x^2+1} = \int_0^{\frac{1}{x_3}} \frac{dx}{x^2+1}$$

From the second equation under (7) we have,

$$x_1 x_2 + x_1 x_3 + x_2 x_3 = 1. \quad \text{From this } x_1 x_2 + (x_1 + x_2) x_3 = 1, \text{ and}$$

$$x_3 = \frac{1 - x_1 x_2}{x_1 + x_2}. \quad \text{Therefore } \frac{1}{x_3} = \frac{x_1 + x_2}{1 - x_1 x_2}$$


Hence (13) becomes,

$$\int_0^{x_1} \frac{dx}{x^2+1} + \int_0^{x_2} \frac{dx}{x^2+1} = \int_0^{\frac{x_1 + x_2}{1 - x_1 x_2}} \frac{dx}{x^2+1}$$

Or, $\text{ARC TANGENT } x_1 + \text{ARC TANGENT } x_2 = \text{ARC TANGENT } \frac{x_1 + x_2}{1 - x_1 x_2}$

Which is, "THE ADDITION THEOREM FOR THE ARC TANGENT".

The values taken for a and b for determining the constant C , $a = 0$, $b = 1$, threw the straight line, used as a variable curve, in the position as shown on the graph below.




SECTION (3)

THE ADDITION THEOREM FOR LEGENDRE'S ELLIPTIC INTEGRAL

OF THE FIRST KIND, $F(k, x) = \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$

- (1) Let $y = \sqrt{(1-x^2)(1-k^2x^2)}$ represent the fixed curve.
 (2) And $y = kx^2 + ax + b$ the variable curve.



$$U(ab) = \int_0^{x_1(ab)} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2(ab)} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_{\frac{1}{k}}^{x_3(ab)} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$$

From (1) and (2) we obtain the cubic,

$2akx^3 + (a^2 + 2bk + k^2 + 1)x^2 + 2abx + b^2 - 1 = 0$ whose roots are the abscissas of the intersections of (1) and (2).

Therefore we have,

(3) $\phi(abx) = 2akx^3 + (a^2 + k^2 + 2bk + 1)x^2 + 2abx + b^2 - 1 = 0$

Proceeding as in SECTIONS (1) and (2), we have,

$$(4) \quad U_a = y_1 x_{1a} + y_2 x_{2a} + y_3 x_{3a}$$

Determining x_a , we have,

$$\phi'_x \cdot x_a + \phi_a = \phi'_x \cdot x_a + (2kx^3 + 2ax^2 + 2bx) = 0$$

$$\text{Therefore } x_a = - \frac{2x(kx^2 + ax + b)}{\phi'_x}$$

$$\text{Where } \phi'_x = 6akx^2 + (a^2 + k^2 + 2bk + 1)2x + 2ab$$

$$\text{Hence, } x_a = - \frac{x(kx^2 + ax + b)}{3akx^2 + (a^2 + k^2 + 2bk + 1)x + ab} = \frac{-xy}{3akx^2 + (a^2 + k^2 + 2bk + 1)x + ab}$$

Substituting in (4) we obtain,

$$(5) \quad U_a = - \left[\frac{1}{y_1} \cdot \frac{x_1 y_1}{3akx_1^2 + (a^2 + k^2 + 2bk + 1)x_1 + ab} + \frac{1}{y_2} \cdot \frac{x_2 y_2}{3akx_2^2 + (a^2 + k^2 + 2bk + 1)x_2 + ab} \right. \\ \left. + \frac{1}{y_3} \cdot \frac{x_3 y_3}{3akx_3^2 + (a^2 + k^2 + 2bk + 1)x_3 + ab} \right] \quad \text{consequently (5) becomes,}$$

$$(6) \quad U_a = - \left[\frac{x_1}{3akx_1^2 + (a^2 + k^2 + 2bk + 1)x_1 + ab} + \frac{x_2}{3akx_2^2 + (a^2 + k^2 + 2bk + 1)x_2 + ab} \right. \\ \left. + \frac{x_3}{3akx_3^2 + (a^2 + k^2 + 2bk + 1)x_3 + ab} \right]$$

(7) By elementary reductions the numerator of U_a becomes,

$$x_1 [9a^2 k^2 x_2^2 x_3^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_2^2 x_3 + x_3^2 x_2) + 3a^2 bk(x_2^2 + x_3^2) \\ + (a^2 + k^2 + 2bk + 1)^2 x_2 x_3 + a^2 b^2 + ab(a^2 + k^2 + 2bk + 1)(x_2 + x_3)] + \\ x_2 [9a^2 k^2 x_1^2 x_3^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2 x_3 + x_3^2 x_1) + 3a^2 bk(x_1^2 + x_3^2) + \\ (a^2 + k^2 + 2bk + 1)^2 x_1 x_3 + a^2 b^2 + ab(a^2 + k^2 + 2bk + 1)(x_1 + x_3)] + \\ x_3 [9a^2 k^2 x_1^2 x_2^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2 x_2 + x_2^2 x_1) + 3a^2 bk(x_1^2 + x_2^2) + \\ (a^2 + k^2 + 2bk + 1)^2 x_1 x_2 + a^2 b^2 + ab(a^2 + k^2 + 2bk + 1)(x_1 + x_2)]$$

Which reduces to,

$$\begin{aligned}
 (8) \quad & 9a^2k^2(x, x_2x_3)(x, x_3 + x, x_2 + x_2x_3) + 6ak(a^2+k^2+2bk+1) \\
 & (x, x_2x_3)(x, + x_2 + x_3) + 3a^2bk(x, x_2^2 + x, x_3^2 + x_2x_1^2 + x_2x_3^2 + x_3x_1^2 + \\
 & x_3x_2^2) + (a^2+k^2+2bk+1)^2(3)(x, x_2x_3) + a^2b^2(x, + x_2 + x_3) + \\
 & 2ab(a^2+k^2+2bk+1)(x, x_2 + x, x_3 + x_2x_3)
 \end{aligned}$$

From (3) we obtain,

$$\begin{aligned}
 (9) \quad & \left\{ \begin{aligned}
 (a) \quad & x_1 + x_2 + x_3 = -\frac{a^2+k^2+2bk+1}{2ak} \\
 (b) \quad & x_1x_2 + x_1x_3 + x_2x_3 = \frac{b}{k} \\
 (c) \quad & x_1x_2x_3 = \frac{1-b^2}{2ak} \\
 & \text{Multiplying (a) by (b) we get,} \\
 (d) \quad & x_1x_2^2 + x_1x_3^2 + x_2x_1^2 + x_2x_3^2 + x_3x_1^2 + x_3x_2^2 = (x_1 + x_2 + x_3) \\
 & (x_1x_2 + x_1x_3 + x_2x_3) - 3(x_1x_2x_3) = \left(-\frac{a^2+k^2+2bk+1}{2ak}\right)\left(\frac{b}{k}\right) - \\
 & (3)\left(\frac{1-b^2}{2ak}\right) = \frac{-a^2b-bk^2+b^2k-b-3k}{2ak^2} \\
 (e) \quad & \text{from (a)} \quad a^2+k^2+2bk+1 = -2ak(x_1 + x_2 + x_3)
 \end{aligned} \right.
 \end{aligned}$$

Substituting these values in (8) we obtain, as the numerator of U_ω ,

$$\begin{aligned}
 & 9a^2k^2\left(\frac{1-b^2}{2ak}\right)\left(\frac{b}{k}\right) + 6ak(a^2+k^2+2bk+1)(x, x_2x_3)(x, + x_2 + x_3) + \\
 & 3a^2bk\left(\frac{-a^2b-bk^2+b^2k-b-3k}{2ak^2}\right) - 6ak(a^2+k^2+2bk+1)(x, x_2x_3) \\
 & (x, + x_2 + x_3) + a^2b^2\left(-\frac{a^2+k^2+2bk+1}{2ak}\right) + 2ab(a^2+k^2+2bk+1)\left(\frac{b}{k}\right) \\
 & = \frac{9ab-9ab^3}{2} + \frac{-3a^3b^2-3ab^2k^2+3ab^3k-3ab^2-9abk}{2k} + \\
 & \frac{-a^3b^2-ab^2k^2-2ab^3k-ab^2}{2k} + \frac{2a^3b^2+2ab^2k^2+4ab^3k+2ab^2}{k} =
 \end{aligned}$$

$$= \frac{9abk - 9ab^3k - 3a^3b^2 - 3ab^2k^2 + 3ab^3k - 3ab^2 - 9abk - a^3b^2 - ab^2k^2 - 2ab^3k}{2k}$$

$$\frac{-ab^2 + 4a^3b^2 + 4ab^2k^2 + 8ab^3k + 4ab^2}{2k} = \frac{0}{2k} = 0$$

$$\text{HENCE } U_a = 0$$

Similarly, for U_b , we have,

$$(10) \quad U_b = y_1 x_{1b} + y_2 x_{2b} + y_3 x_{3b}$$

$$\text{and since } \phi'_x \cdot x_b + \phi_b = \phi'_x \cdot x_b + (2kx^2 + 2ax + 2b) = 0$$

$$\text{Therefore } x_b = - \frac{2kx^2 + 2ax + 2b}{6akx^2 + (a^2 + k^2 + 2bk + 1)2x + 2ab} =$$

$$- \frac{kx^2 + ax + b}{3akx^2 + (a^2 + k^2 + 2bk + 1)x + ab} = \frac{-y}{3akx^2 + (a^2 + k^2 + 2bk + 1)x + ab}$$

Substituting in equation (6), we get,

$$(11) \quad U_b = - \left[\frac{1}{y_1} \cdot \frac{y_1}{3akx_1^2 + (a^2 + k^2 + 2bk + 1)x_1 + ab} + \frac{1}{y_2} \cdot \frac{y_2}{3akx_2^2 + (a^2 + k^2 + 2bk + 1)x_2 + ab} + \frac{1}{y_3} \cdot \frac{y_3}{3akx_3^2 + (a^2 + k^2 + 2bk + 1)x_3 + ab} \right]$$

By elementary reductions, we obtain for the numerator

of U_b ,

$$\begin{aligned} & 9a^2k^2x_2^2x_3^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_2^2x_3 + x_3^2x_2) + \\ & 3a^2bk(x_2^2 + x_3^2) + (a^2 + k^2 + 2bk + 1)^2x_2x_3 + ab(a^2 + k^2 + 2bk + 1)(x_2 + x_3) + \\ & a^2b^2 + 9a^2k^2x_1^2x_3^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2x_3 + x_3^2x_1) + 3a^2bk(x_1^2 + x_3^2) \\ & + (a^2 + k^2 + 2bk + 1)^2x_1x_3 + ab(a^2 + k^2 + 2bk + 1)(x_1 + x_3) + a^2b^2 + \\ & 9a^2k^2x_1^2x_2^2 + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2x_2 + x_2^2x_1) + 3a^2bk(x_1^2 + x_2^2) + \\ & (a^2 + k^2 + 2bk + 1)^2x_1x_2 + ab(a^2 + k^2 + 2bk + 1)(x_1 + x_2) + a^2b^2 = \end{aligned}$$

$$9a^2k^2(x_1^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2) + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2x_2 + x_2^2x_1 +$$

$$x_1^2 x_3 + x_3^2 x_1 + x_2^2 x_3 + x_3^2 x_2) + 3a^2 bk(x_1^2 + x_2^2 + x_3^2 + x_2^2 + x_3^2) + (a^2 + k^2 + 2bk + 1)^2(x, x_2 + x, x_3 + x_2 x_3) + ab(a^2 + k^2 + 2bk + 1)(x, + x_2 + x_1 + x_3 + x_2 + x_3) + 3a^2 b^2 =$$

$$(12) \quad 9a^2 k^2(x_1^2 x_2^2 + x_1^2 x_3^2 + x_2^2 x_3^2) + 3ak(a^2 + k^2 + 2bk + 1)(x_1^2 x_2 + x_2^2 x_1 + x_1^2 x_3 + x_3^2 x_1 + x_2^2 x_3 + x_3^2 x_2) + 6a^2 bk(x_1^2 + x_2^2 + x_3^2) + (a^2 + k^2 + 2bk + 1)^2(x, x_2 + x, x_3 + x_2 x_3) + 2ab(a^2 + k^2 + 2bk + 1)(x, + x_2 + x_3) + 3a^2 b^2$$

From (3) we have,

$$(13) \quad \begin{cases} x_1 + x_2 + x_3 = -\frac{a^2 + k^2 + 2bk + 1}{2ak} \\ x_1 x_2 + x_1 x_3 + x_2 x_3 = \frac{b}{k} \\ x_1 x_2 x_3 = \frac{1 - b^2}{2ak} \\ x_1^2 + x_2^2 + x_3^2 = (x_1 + x_2 + x_3)^2 - 2(x_1 x_2 + x_1 x_3 + x_2 x_3) \\ x_1^2 x_2^2 + x_1^2 x_3^2 + x_2^2 x_3^2 = (x_1 x_2 + x_1 x_3 + x_2 x_3)^2 - 2(x_1 x_2 x_3)(x_1 + x_2 + x_3) \\ x_1^2 x_2 + x_2^2 x_1 + x_1^2 x_3 + x_3^2 x_1 + x_2^2 x_3 + x_3^2 x_2 = (x_1 + x_2 + x_3)(x_1 x_2 + x_1 x_3 + x_2 x_3) - 3(x_1 x_2 x_3) \end{cases}$$

Substituting these values in (12) we obtain, as numerator of U_b ,

$$9a^2 k^2 \left[(x_1 x_2 + x_1 x_3 + x_2 x_3)^2 - 2(x_1 x_2 x_3)(x_1 + x_2 + x_3) \right] + 3ak(a^2 + k^2 + 2bk + 1) \left[(x_1 + x_2 + x_3)(x_1 x_2 + x_1 x_3 + x_2 x_3) - 3(x_1 x_2 x_3) \right] + 6a^2 bk \left[(x_1 + x_2 + x_3)^2 - 2(x_1 x_2 + x_1 x_3 + x_2 x_3) \right] + (a^2 + k^2 + 2bk + 1)^2(x, x_2 + x, x_3 + x_2 x_3) + 2ab(a^2 + k^2 + 2bk + 1)(x, + x_2 + x_3) + 3a^2 b^2 =$$

$$9a^2 k^2 \left[\frac{b^2}{k^2} - 2 \left(\frac{1 - b^2}{2ak} \right) \left(-\frac{a^2 + k^2 + 2bk + 1}{2ak} \right) \right] +$$

$$\begin{aligned}
& 3ak(a^2+k^2+2bk+1) \left[\left(-\frac{a^2+k^2+2bk+1}{2ak} \right) \left(\frac{b}{k} \right) - 3 \left(\frac{1-b^2}{2ak} \right) \right] + \\
& 6a^2bk \left[\left(-\frac{a^2+k^2+2bk+1}{2ak} \right)^2 - \frac{2b}{k} \right] + \cancel{(a^2+k^2+2bk+1)^2 \left(\frac{b}{k} \right)} + \\
& \cancel{2ab(a^2+k^2+2bk+1) \left(-\frac{a^2+k^2+2bk+1}{2ak} \right)} + 3a^2b^2
\end{aligned}$$

The quantities crossed in red are equal and opposite in sign and vanish. Partially multiplying the remaining terms out, we get, as the numerator of U_b ,

$$\begin{aligned}
& 9a^2b^2 + \frac{9(1-b^2)(a^2+k^2+2bk+1)}{2} - \frac{3b(a^2+k^2+2bk+1)^2}{2k} - \\
& \frac{9(1-b^2)(a^2+k^2+2bk+1)}{2} + \frac{3b(a^2+k^2+2bk+1)^2}{2k} - 12a^2b^2 + 3a^2b^2 = 0
\end{aligned}$$

$$\text{Therefore } U_b = 0$$

HENCE $U_a = 0$, $U_b = 0$ and we have therefore,

$$U(ab) = \text{a constant} = C$$

We have, then,

$$(14) \int_0^{x_1} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_{\frac{1}{k}}^{x_3} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = C$$

Determination of C

If, in (3), we put $b = 1$ and $a = 0$, it reduces to

$(k+1)^2 x^2 = 0$ or $x^2 = 0$, i.e. two roots of the cubic are 0 and the third increases without bound, giving $x_1 = x_2 = 0, x_3 = \infty$

Substituting in (14) we have,

$$(15) \int_0^0 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^0 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_{\frac{1}{k}}^{\infty} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = C$$

The first two integrals in (15) vanish identically, and putting $x = \frac{1}{kt}$, $dx = -\frac{dt}{kt^2}$, the third integral becomes,

$$-\int_1^{t_3} \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}} = C, \quad \text{As } x \rightarrow \infty, \quad t_3 = \frac{1}{kx_3} \rightarrow \frac{1}{\infty} = 0$$

Consequently, $\int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}} = C$, and (14) becomes,

$$(16) \int_0^{x_1} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} - \int_1^{t_3} \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}} =$$

$$\int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}.$$

$$\text{But, } \int_1^{t_3} f(t)dt = \int_1^0 f(t)dt + \int_0^{t_3} f(t)dt = -\int_0^1 f(t)dt + \int_0^{t_3} f(t)dt$$

Therefore substitution in (16) gives,

$$\int_0^{x_1} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = \int_0^{t_3} \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}$$

Replacing t_3 by $\frac{1}{kx_3}$, we get,

$$(17) \int_0^{x_1} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = \int_0^{\frac{1}{kx_3}} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$$

From equation (3),

$$x_1 x_2 x_3 = \frac{1-b^2}{2ak}$$

Therefore $kx_1 x_2 = \frac{1-b^2}{2ax_3}$, and $k^2 x_1^2 x_2^2 = \frac{(1-b^2)^2}{4a^2 x_3^2}$

By Composition, $1 - k^2 x_1^2 x_2^2 = 1 - \frac{(1-b^2)^2}{4a^2 x_3^2} = \frac{4a^2 x_3^2 - (1-b^2)^2}{4a^2 x_3^2}$

Or, $4a^2 x_3^2 = \frac{4a^2 x_3^2 - (1-b^2)^2}{1 - k^2 x_1^2 x_2^2}$, Therefore, we have,

$$(18) \quad x_3 = \frac{\frac{4a^2 x_3^2 - (1-b^2)^2}{4a^2 x_3}}{1 - k^2 x_1^2 x_2^2} = \frac{\frac{4a^2 k x_3^2 - k(1-b^2)^2}{4a^2 k x_3}}{1 - k^2 x_1^2 x_2^2}$$

By multiplying both sides of (18) by $\frac{1}{kx_3^2}$, we obtain,

$$(19) \quad \frac{1}{kx_3} = \frac{\frac{4a^2 k x_3^2 - k(1-b^2)^2}{4a^2 k^2 x_3^3}}{1 - k^2 x_1^2 x_2^2}$$

By separating the numerator of (19) in two parts, and adding and subtracting $\frac{b^2}{kx_3}$, we get,

$$(20) \quad \frac{1}{kx_3} = \frac{\frac{1}{kx_3} - \frac{k(1-b^2)^2}{4a^2 k^2 x_3^3} + \frac{b^2}{kx_3} - \frac{b^2}{kx_3}}{1 - k^2 x_1^2 x_2^2}$$

If, to the numerator of (20), we add and subtract

$\frac{b(1-b^2)}{2akx_3^2}$, it becomes,

$$(21) \quad \frac{1}{kx_3} = \frac{\frac{b(1-b^2)}{2akx_3^2} - \frac{k(1-b^2)^2}{4a^2 k^2 x_3^3} + \frac{1}{kx_3} - \frac{b^2}{kx_3} + \frac{b^2}{kx_3} - \frac{b(1-b^2)}{2akx_3^2}}{1 - k^2 x_1^2 x_2^2}$$

Combining the third and fourth terms of the numerator of

(21) and multiplying and dividing the result by $2a$, we get,

$$(22) \quad \frac{1}{kx_3} = \frac{\frac{b(1-b^2)}{2akx_3^2} - \frac{k(1-b^2)^2}{4a^2k^2x_3^3} + \frac{2a(1-b^2)}{2akx_3} + \frac{b^2}{kx_3} - \frac{b(1-b^2)}{2akx_3^2}}{1 - k^2x_1^2x_2^2}$$

which may be written,

$$(23) \quad \frac{1}{kx_3} = \frac{\frac{b\left(\frac{1-b^2}{2akx_3}\right) - k\left(\frac{(1-b^2)^2}{4a^2k^2x_3^2}\right)}{x_3} + 2a\left(\frac{1-b^2}{2akx_3}\right) + \frac{b\left(\frac{b}{k} - \frac{1-b^2}{2akx_3}\right)}{x_3}}{1 - k^2x_1^2x_2^2}$$

From $x_1x_2x_3 = \frac{1-b^2}{2ak}$ we get,

$$(24) \quad \frac{1-b^2}{2akx_3} = x_1x_2 \quad \text{and} \quad (25) \quad \frac{(1-b^2)^2}{4a^2k^2x_3^2} = x_1^2x_2^2$$

Substituting these values in (23) we obtain,

$$(26) \quad \frac{1}{kx_3} = \frac{\frac{bx_1x_2 - kx_1^2x_2^2}{x_3} + 2ax_1x_2 + \frac{b\left(\frac{b}{k} - x_1x_2\right)}{x_3}}{1 - k^2x_1^2x_2^2}$$

and, changing form to obtain the quotient $\frac{\frac{b}{k} - x_1x_2}{x_3}$,

(26) becomes,

$$(27) \quad \frac{1}{kx_3} = \frac{kx_1x_2\left(\frac{\frac{b}{k} - x_1x_2}{x_3}\right) + 2ax_1x_2 + b\left(\frac{\frac{b}{k} - x_1x_2}{x_3}\right)}{1 - k^2x_1^2x_2^2}$$

From (3), $x_1x_2 + x_1x_3 + x_2x_3 = \frac{b}{k}$

$$x_1 x_2 + x_3 (x_1 + x_2) = \frac{b}{k}, \text{ therefore } x_3 (x_1 + x_2) = \frac{b}{k} - x_1 x_2$$

$$(28) \quad \text{Hence, } \frac{\frac{b}{k} - x_1 x_2}{x_3} = x_1 + x_2$$

Using (28), (27) becomes,

$$\begin{aligned} \frac{1}{kx_3} &= \frac{kx_1 x_2 (x_1 + x_2) + 2ax_1 x_2 + b(x_1 + x_2)}{1 - k^2 x_1^2 x_2^2} \\ &= \frac{kx_1^2 x_2 + kx_1 x_2^2 + ax_1 x_2 + ax_1 x_2 + bx_1 + bx_2}{1 - k^2 x_1^2 x_2^2} \\ &= \frac{(kx_1 x_2^2 + ax_1 x_2 + bx_1) + (kx_1^2 x_2 + ax_1 x_2 + bx_2)}{1 - k^2 x_1^2 x_2^2} \end{aligned}$$

$$(29) \quad \frac{1}{kx_3} = \frac{x_1 (kx_2^2 + ax_2 + b) + x_2 (kx_1^2 + ax_1 + b)}{1 - k^2 x_1^2 x_2^2}$$

But, since x_1 and x_2 are roots of,

$$kx^2 + ax + b - \sqrt{(1-x^2)(1-k^2x^2)} = 0,$$

$$kx_1^2 + ax_1 + b = \sqrt{(1-x_1^2)(1-k^2x_1^2)}$$

$$\text{and, } kx_2^2 + ax_2 + b = \sqrt{(1-x_2^2)(1-k^2x_2^2)}$$

therefore, substituting in (29), we get,

$$(30) \quad \frac{1}{kx_3} = x_3' = \frac{x_1 \sqrt{(1-x_2^2)(1-k^2x_2^2)} + x_2 \sqrt{(1-x_1^2)(1-k^2x_1^2)}}{1 - k^2 x_1^2 x_2^2}$$

Consequently, (17) becomes,

$$\int_0^{x_1} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \int_0^{x_2} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = \int_0^{x'_3} \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$$

or, $F(k, x_1) + F(k, x_2) = F(k, x'_3)$ where x'_3 has the value given in (30).

This is, "THE ADDITION THEOREM FOR THE LEGENDRE ELLIPTIC INTEGRAL OF THE FIRST KIND .", $F(k, x)$.

For the special values of the cubic, $a = 0$ and $b = 1$, the variable curve takes the position on the graph as shown below:

